

Automorphism Groups of Simple Moufang Loops over Perfect Fields

BY GÁBOR P. NAGY *

SZTE Bolyai Institute

Aradi vértanúk tere 1, H-6720 Szeged, Hungary

e-mail: nagy@math.u-szeged.hu

PETR VOJTĚCHOVSKÝ †

Department of Mathematics, Iowa State University

400 Carver Hall, Ames, Iowa, 50011, USA

e-mail: petr@iastate.edu

February 2, 2008

Abstract

Let F be a perfect field and $M(F)$ the nonassociative simple Moufang loop consisting of the units in the (unique) split octonion algebra $O(F)$ modulo the center. Then $\text{Aut}(M(F))$ is equal to $G_2(F) \rtimes \text{Aut}(F)$. In particular, every automorphism of $M(F)$ is induced by a semilinear automorphism of $O(F)$. The proof combines results and methods from geometrical loop theory, groups of Lie type and composition algebras; its gist being an identification of the automorphism group of a Moufang loop with a subgroup of the automorphism group of the associated group with triality.

1 Introduction

As we hope to attract the attention of both group- and loop-theorists, we take the risk of being trivial at times and introduce most of the background material carefully, although briefly. We refer the reader to [11], [10], [3] and [7] for a more systematic exposition.

A groupoid Q is a *quasigroup* if the equation $xy = z$ has a unique solution in Q whenever two of the three elements $x, y, z \in Q$ are known. A *loop* is a quasigroup with a neutral element, denoted by 1 in the sequel. *Moufang loop*

*Supported by the “János Bolyai Fellowship” of the Hungarian Academy of Sciences, and by the grants OTKA T029849 and FKFP 0063/2001.

†Partially supported by the Grant Agency of Charles University, grant no. 269/2001/B-MAT/MFF, and by research assistantship at Iowa State University.

is a loop satisfying one of the (equivalent) *Moufang identities*, for instance the identity $((xy)x)z = x(y(xz))$. The *multiplication group* $\text{Mlt}(L)$ of a loop L is the group generated by all left and right translations $x \mapsto ax$, $x \mapsto xa$, where $a \in L$.

Let C be a vector space over a field F , and $N : C \rightarrow F$ a nondegenerate quadratic form. Define multiplication $\cdot$ on C so that $(C, +, \cdot)$ becomes a not necessarily associative ring. Then $C = (C, N)$ is a *composition algebra* if $N(u \cdot v) = N(u) \cdot N(v)$ holds for every $u, v \in C$. Composition algebras exist only in dimensions 1, 2, 4 and 8, and we speak of an *octonion algebra* when $\dim C = 8$. A composition algebra is called *split* when it has nontrivial zero divisors. By [11, Theorem 1.8.1], there is a unique split octonion algebra $O(F)$ over any field F .

Write $O(F)^*$ for the set of all elements of unit norm in $O(F)$, and let $M(F)$ be the quotient of $O(F)^*$ by its center $Z(O(F)^*) = \{\pm 1\}$. Since every composition algebra satisfies all Moufang identities, both $O(F)^*$ and $M(F)$ are Moufang loops. Paige proved [9] that $M(F)$ is nonassociative and simple (as a loop). Liebeck [8] used the classification of finite simple groups to conclude that there are no other nonassociative finite simple Moufang loops besides $M(F)$, F finite.

Liebeck's proof relies heavily on results of Doro [4], that relate Moufang loops to groups with triality. Before we define these groups, allow us to say a few words about the (standard) notation. Let G be a group. Working in $G \rtimes \text{Aut}(G)$, when $g \in G$ and $\alpha \in \text{Aut}(G)$, we write g^α for the image of g under α , and $[g, \alpha]$ for $g^{-1}g^\alpha$. Appealing to this convention, we say that α *centralizes* g if $g^\alpha = g$. Now, the pair (G, S) is said to be a *group with triality* if $S \leq \text{Aut}(G)$, $S = \langle \sigma, \rho \rangle \cong S_3$, σ is an involution, ρ is of order 3, $G = [G, S]$, $Z(GS) = \{1\}$, and the triality equation

$$[g, \sigma][g, \sigma]^\rho[g, \sigma]^{\rho^2} = 1$$

holds for every $g \in G$.

We now turn to geometrical loop theory. A *3-net* is an incidence structure $\mathcal{N} = (\mathcal{P}, \mathcal{L})$ with point set $\mathcal{P}$ and line set $\mathcal{L}$, where $\mathcal{L}$ is a disjoint union of 3 classes $\mathcal{L}_i$ ($i = 1, 2, 3$) such that two distinct lines from the same class have no point in common, and any two lines from distinct classes intersect in exactly one point. A line from the class $\mathcal{L}_i$ is usually referred to as an *i-line*. A permutation on $\mathcal{P}$ is a *collineation* of $\mathcal{N}$ if it maps lines to lines. We speak of a *direction preserving* collineation if the line classes $\mathcal{L}_i$ are invariant under the induced permutation of lines.

There is a canonical correspondence between loops and 3-nets. Any loop L determines a 3-net when we let $\mathcal{P} = L \times L$, $\mathcal{L}_1 = \{(c, y) | y \in L\} | c \in L$, $\mathcal{L}_2 = \{(x, c) | x \in L\} | c \in L$, $\mathcal{L}_3 = \{(x, y) | x, y \in L, xy = c\} | c \in L$. Conversely, given a 3-net $\mathcal{N} = (\mathcal{P}, \mathcal{L})$ and the origin $1 \in \mathcal{P}$, we can introduce multiplication on the 1-line ℓ through 1 that turns ℓ into a loop, called the *coordinate loop* of $\mathcal{N}$. Since the details of this construction are not essential for what follows, we omit them.

Let $\mathcal{N}$ be a 3-net and $\ell_i \in \mathcal{L}_i$, for some i . We define a certain permutation σ_{ℓ_i} on the point set $\mathcal{P}$ (cf. Figure 1). For $P \in \mathcal{P}$, let a_j and a_k be the lines through




Figure 1: The Bol reflection with axis ℓ_1

P such that $a_j \in \mathcal{L}_j$, $a_k \in \mathcal{L}_k$, and $\{i, j, k\} = \{1, 2, 3\}$. Then there are unique intersection points $Q_j = a_j \cap \ell_i$, $Q_k = a_k \cap \ell_i$. We define $\sigma_{\ell_i}(P) = b_j \cap b_k$, where b_j is the unique j -line through Q_k , and b_k the unique k -line through Q_j . The permutation σ_{ℓ_i} is clearly an involution satisfying $\sigma_{\ell_i}(\mathcal{L}_j) = \mathcal{L}_k$, $\sigma_{\ell_i}(\mathcal{L}_k) = \mathcal{L}_j$. If it happens to be the case that σ_{ℓ_i} is a collineation, we call it the *Bol reflection with axis ℓ_i* .

It is clear that for any collineation γ of $\mathcal{N}$ and any line ℓ we have $\sigma_{\gamma(\ell)} = \gamma \sigma_{\ell} \gamma^{-1}$. Hence the set of Bol reflections of $\mathcal{N}$ is invariant under conjugations by elements of the collineation group $\text{Coll}(\mathcal{N})$ of $\mathcal{N}$. A 3-net $\mathcal{N}$ is called a *Moufang 3-net* if σ_{ℓ} is a Bol reflection for every line ℓ . Bol proved that $\mathcal{N}$ is a Moufang 3-net if and only if all coordinate loops of $\mathcal{N}$ are Moufang (cf. [2, p. 120]).

We are now coming to the crucial idea of this paper. For a Moufang 3-net $\mathcal{N}$ with origin 1, denote by ℓ_i ($i = 1, 2, 3$) the three lines through 1. As in [7], we write Γ_0 for the subgroup of $\text{Coll}(\mathcal{N})$ generated by all Bol reflections of $\mathcal{N}$, and Γ for the direction preserving part of Γ_0 . Also, let S be the subgroup generated by σ_{ℓ_1} , σ_{ℓ_2} and σ_{ℓ_3} . According to [7], Γ is a normal subgroup of index 6 in Γ_0 , $\Gamma_0 = \Gamma S$, and (Γ, S) is a group with triality. (Here, S is understood as a subgroup of $\text{Aut}(\Gamma)$ by identifying $\sigma \in S$ with the map $\tau \mapsto \sigma \tau \sigma^{-1}$.) We will always fix $\sigma = \sigma_{\ell_1}$ and $\rho = \sigma_{\ell_1} \sigma_{\ell_2}$ in such a situation, to obtain $S = \langle \sigma, \rho \rangle$ as in the definition of a group with triality.

2 The Automorphisms

Let C be a composition algebra over F . A map $\alpha : C \rightarrow C$ is a *linear automorphism* (resp. *semilinear automorphism*) of C if it is a bijective F -linear (resp. F -semilinear) map preserving the multiplication, i.e., satisfying $\alpha(uv) = \alpha(u)\alpha(v)$ for every $u, v \in C$. It is well known that the group of linear automorphisms of

$O(F)$ is isomorphic to the Chevalley group $G_2(F)$, cf. [5, Section 3], [11, Chapter 2]. The group of semilinear automorphisms of $O(F)$ is therefore isomorphic to $G_2(F) \rtimes \text{Aut}(F)$.

Since every linear automorphism of a composition algebra is an isometry [11, Section 1.7], it induces an automorphisms of the loop $M(F)$. By [12, Theorem 3.3], every element of $O(F)$ is a sum of two elements of norm one. Consequently, $\text{Aut}(O(F)) \leq \text{Aut}(M(F))$.

An automorphism $f \in \text{Aut}(M(F))$ will be called *(semi)linear* if it is induced by a (semi)linear automorphism of $O(F)$. By considering extensions of automorphisms of $M(F)$, it was proved in [12] that $\text{Aut}(M(\mathbb{F}_2))$ is isomorphic to $G_2(\mathbb{F}_2)$, where $\mathbb{F}_2$ is the two-element field. The aim of this paper is to generalize this result (although using different techniques) and prove that every automorphism of $\text{Aut}(M(F))$ is semilinear, provided F is perfect. We reach this aim by identifying $\text{Aut}(M(F))$ with a certain subgroup of the automorphism group of the group with triality associated with $M(F)$.

To begin with, we recall the geometrical characterization of automorphisms of a loop.

Lemma 2.1 (Theorem 10.2 [1]) *Let L be a loop and $\mathcal{N}$ its associated 3-net. Any direction preserving collineation which fixes the origin of $\mathcal{N}$ is of the form $(x, y) \mapsto (x^\alpha, y^\alpha)$ for some $\alpha \in \text{Aut}(L)$. Conversely, the map $\alpha : L \rightarrow L$ is an automorphism of L if and only if $(x, y) \mapsto (x^\alpha, y^\alpha)$ is a direction preserving collineation of $\mathcal{N}$.*

We will denote the map $(x, y) \mapsto (x^\alpha, y^\alpha)$ by φ_α .

By [7, Propositions 3.3 and 3.4], $\mathcal{N}$ is embedded in $\Gamma_0 = \Gamma S$ as follows. The lines of $\mathcal{N}$ correspond to the conjugacy classes of σ in Γ_0 , two lines are parallel if and only if the corresponding involutions are Γ -conjugate, and three pairwise non-parallel lines have a point in common if and only if they generate a subgroup isomorphic to S_3 . In particular, the three lines through the origin of $\mathcal{N}$ correspond to the three involutions of S .

As the set of Bol reflections of $\mathcal{N}$ is invariant under conjugations by collineations, every element $\varphi \in \text{Coll}(\mathcal{N})$ normalizes the group Γ and induces an automorphism $\widehat{\varphi}$ of Γ . It is not difficult to see that φ fixes the three lines through the origin of $\mathcal{N}$ if and only if $\widehat{\varphi}$ centralizes (the involutions of) S .

Proposition 2.2 *Let L be a Moufang loop and $\mathcal{N}$ its associated 3-net. Let Γ_0 be the group of collineations generated by the Bol reflections of $\mathcal{N}$, Γ the direction preserving part of Γ_0 , and $S \cong S_3$ the group generated by the Bol reflections whose axis contains the origin of $\mathcal{N}$. Then $\text{Aut}(L) \cong C_{\text{Aut}(\Gamma)}(S)$.*

Proof: Pick $\alpha \in \text{Aut}(L)$, and let $\widehat{\varphi}_\alpha$ be the automorphism of Γ induced by the collineation φ_α . As φ_α fixes the three lines through the origin, $\widehat{\varphi}_\alpha$ belongs to $C_{\text{Aut}(\Gamma)}(S)$.

Conversely, an element $\psi \in C_{\text{Aut}(\Gamma)}(S)$ normalizes the conjugacy class of σ in ΓS and preserves the incidence structure defined by the embedding of $\mathcal{N}$. This means that $\psi = \widehat{\varphi}$ for some collineation $\varphi \in \text{Coll}(\mathcal{N})$. Now, ψ centralizes

S , therefore φ fixes the three lines through the origin. Thus φ must be direction preserving, and there is $\alpha \in \text{Aut}(L)$ such that $\varphi = \varphi_\alpha$, by Lemma 2.1.

It remains to add the last ingredient—groups of Lie type.

Theorem 2.3 *Let F be a perfect field. Then the automorphism group of the nonassociative simple Moufang loop $M(F)$ constructed over F is isomorphic to the semidirect product $G_2(F) \rtimes \text{Aut}(F)$. Every automorphism of $M(F)$ is induced by a semilinear automorphism of the split octonion algebra $O(F)$.*

Proof: We fix a perfect field F , and assume that all simple Moufang loops and Lie groups mentioned below are constructed over F .

The group with triality associated with M turns out to be its multiplicative group $\text{Mlt}(M) \cong D_4$, and the graph automorphisms of D_4 are exactly the triality automorphisms of M (cf. [5], [4]). To be more precise, Freudenthal proved this for the reals and Doro for finite fields, however they based their arguments only on the root system and parabolic subgroups, and that is why their result is valid over any field.

By [5], $C_{D_4}(\sigma) = B_3$, and by [8, Lemmas 4.9, 4.10 and 4.3], $C_{D_4}(\rho) = G_2$. As $G_2 < B_3$, by [6, p. 28], we have $C_{D_4}(S_3) = G_2$.

Since F is perfect, $\text{Aut}(D_4)$ is isomorphic to $\Delta \rtimes (\text{Aut}(F) \times S_3)$, by a result of Steinberg (cf. [3, Chapter 12]). Here, Δ is the group of the inner and diagonal automorphisms of D_4 , and S_3 is the group of graph automorphisms of D_4 . When $\text{char } F = 2$ then no diagonal automorphisms exist, and $\Delta = \text{Inn}(D_4)$. When $\text{char } F \neq 2$ then S_3 acts faithfully on $\Delta/\text{Inn}(D_4) \cong C_2 \times C_2$. Hence, in any case, $C_\Delta(S_3) = C_{D_4}(S_3)$. Moreover, for the field and graph automorphisms commute, we have $C_{\text{Aut}(D_4)}(S_3) = C_{D_4}(S_3) \rtimes \text{Aut}(F)$.

We have proved $\text{Aut}(M) \cong G_2 \rtimes \text{Aut}(F)$. The last statement follows from the fact that the group of linear automorphisms of the split octonion algebra is isomorphic to G_2 .

One of the open questions in loop theory is to decide which groups can be obtained as multiplication groups of loops. Thinking along these lines we ask: *Which groups can be obtained as automorphism groups of loops?* Theorem 2.3 yields a partial answer. Namely, every Lie group of type G_2 over a perfect field can be obtained in this way.

Finally, the former author asked the latter one at the Loops '99 conference whether $\text{Aut}(M(F))$ is simple when F is finite. We now know that this happens if and only if F is a finite prime field of odd characteristic.

References

- [1] A. BARLOTTI AND K. STRAMBACH. The geometry of binary systems. *Advances Math.* **49** (1983), 1–105.
- [2] R.H. BRUCK. *A survey of binary systems*. (Springer-Verlag, Berlin, 1958).
- [3] R.W. CARTER. *Simple groups of Lie type*. (Wiley Interscience, 1972).

- [4] S. DORO. Simple Moufang loops. *Math. Proc. Cambridge Philos. Soc.* **83** (1978), 377–392.
- [5] H. FREUDENTHAL. Oktaven, Ausnahmegruppen und Oktavengeometrie. *Geometriae Dedicata* **19** (1985), 1–63.
- [6] D. GORENSTEIN, R. LYONS and R. SOLOMON. *The classification of the finite simple groups*. No. 3. Part I, Mathematical Surveys and Monographs **40**(3) (Providence, R.I., AMS, 1998).
- [7] J.I. HALL and G. P. NAGY. On Moufang 3-nets and groups with triality. *Acta Sci. Math. (Szeged)* **67** (2001), 675–685.
- [8] M.W. LIEBECK. The classification of finite simple Moufang loops. *Math. Proc. Cambridge Philos. Soc.* **102** (1987), 33–47.
- [9] L.J. PAIGE. A class of simple Moufang loops. *Proc. Amer. Math. Soc.* **7** (1956), 471–482.
- [10] H.O. PFLUGFELDER. *Quasigroups and Loops: Introduction*, Sigma series in pure mathematics; **7** (Heldermann Verlag Berlin, 1990).
- [11] T.A. SPRINGER and F. D. VELDKAMP, *Octonions, Jordan Algebras, and Exceptional Groups*, Springer Monographs in Mathematics (Springer Verlag, 2000).
- [12] P. VOJTĚCHOVSKÝ. Finite simple Moufang loops. PhD Thesis, Iowa State University, 2001.